

fielder Sulphate

**Contains: 825 grammes per litre Sulphur Trioxide (62% S₀₃ w/w)
330 grammes per litre Sulphur (26% S w/w)
140 grammes per litre Ammoniacal Nitrogen (11% N w/w)**

An inorganic liquid formulation of sulphur formulated in sulphate form. Elemental sulphur is not readily available to plant as a nutrient source and needs to be first oxidised to the sulphate salt before significant uptake is possible. Fielder Sulphate is purposely formulated for immediate leaf uptake and maximum plant availability. The purpose of significant level of nitrogen also aids protein formulation improving plant health growth and quality. Fielder Sulphate contains its own surfactants to improve tank mixing.

RECOMMENDATIONS FOR USE:

Maintenance - Apply 5 l/ha (3.5 pints/acre) as soon as there is sufficient leaf area to absorb the spray.
Deficiency or severe stress - Apply 10 l/ha (7 pints/acre) as soon as there is sufficient leaf area or as soon as symptoms appear. Repeat every 14 days if required.

CROP RECOMMENDATIONS:

Fielder Sulphate is recommended for use on most arable, horticultural & fruit

WATER RATE:

Conventional sprayer : 220 litres/ha (20 galls/acre)
Low volume sprayer : 55 litres/ha (5 galls/acre)

TANK MIXING:

Fielder Sulphate is physically and chemically compatible with most pesticides. Check by mixing small amount in the correct ratio adding the product to the water followed by pesticide. Crops can be susceptible to scorch from mixtures of manganese and pesticides applied during periods of rapid growth or stress. Always add Fielder Sulphate to spray tank and then add pesticide.

PRECAUTIONS:

For the best results apply when crop actively growing. Ensure spray equipment is free of residues from previous use. Shake product container thoroughly before adding to the sprayer. Mix for five minutes before spraying. Do not allow mixture to stand idle at any time. Avoid spraying during very high temperatures & bright sunlight. Keep out of reach of children. Avoid contact with skin, eyes and clothing. Wash exposed area with water. Prevent livestock access to treated area for 14 days. Protect from frost. Contents may be harmful to marine life; do not contaminate waterways, ditches, lakes or ponds with containers. Rinse residues into spray tank and dispose of containers safely.

CONDITIONS OF SUPPLY:

All goods supplied by us are of a high quality and we believe them to be suitable, but as we cannot exercise control over their storage, handling, mixing or use or of the weather conditions before, during or after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

**Contents:
10 Litres e**

fielder

The Paddocks - Longden - Shrewsbury - Shropshire - SY5 8EX

Tel: 01743 860924 - Fax: 01743 860977